

A class of hypergraphs that generalizes chordal graphs

Eric Emtander*

Abstract

In this paper we introduce a class of hypergraphs that we call chordal. We also extend the definition of triangulated hypergraphs, given in [9], so that a triangulated hypergraph, according to our definition, is a natural generalization of a chordal (rigid circuit) graph. In [7], Fröberg shows that the chordal graphs corresponds to graph algebras, $R/I(\mathcal{G})$, with linear resolutions. We extend Fröberg's method and show that the hypergraph algebras of generalized chordal hypergraphs, a class of hypergraphs that includes the chordal hypergraphs, have linear resolutions. The definitions we give, yield a natural higher dimensional version of the well known flag property of simplicial complexes. We obtain what we call d -flag complexes.

1 Introduction and preliminaries

Let $\mathcal{X}$ be a finite set and $\mathcal{E} = \{E_1, \dots, E_s\}$ a finite collection of non empty subsets of $\mathcal{X}$. The pair $\mathcal{H} = (\mathcal{X}, \mathcal{E})$ is called a **hypergraph**. The elements of $\mathcal{X}$ and $\mathcal{E}$, respectively, are called the **vertices** and the **edges**, respectively, of the hypergraph. If we want to specify what hypergraph we consider, we may write $\mathcal{X}(\mathcal{H})$ and $\mathcal{E}(\mathcal{H})$ for the vertices and edges respectively. A hypergraph is called **simple** if: (1) $|E_i| \geq 2$ for all $i = 1, \dots, s$ and (2) $E_j \subseteq E_i$ implies $i = j$. If the cardinality of $\mathcal{X}$ is n we often just use the set $[n] = \{1, 2, \dots, n\}$ instead of $\mathcal{X}$.

Let $\mathcal{H}$ be a hypergraph. A **subhypergraph** $\mathcal{K}$ of $\mathcal{H}$ is a hypergraph such that $\mathcal{X}(\mathcal{K}) \subseteq \mathcal{X}(\mathcal{H})$, and $\mathcal{E}(\mathcal{K}) \subseteq \mathcal{E}(\mathcal{H})$. If $\mathcal{Y} \subseteq \mathcal{X}$, the **induced hypergraph on $\mathcal{Y}$** , $\mathcal{H}_{\mathcal{Y}}$, is the subhypergraph with $\mathcal{X}(\mathcal{H}_{\mathcal{Y}}) = \mathcal{Y}$ and with $\mathcal{E}(\mathcal{H}_{\mathcal{Y}})$ consisting of the edges of $\mathcal{H}$ that lie entirely in $\mathcal{Y}$. A hypergraph $\mathcal{H}$ is said to be **d -uniform** if $|E_i| = d$ for every edge $E_i \in \mathcal{E}(\mathcal{H})$. Note that a simple 2-uniform hypergraph is just an ordinary simple graph.

Throughout the paper we denote by R the polynomial ring $k[x_1, \dots, x_n]$ over some field k , where n is the number of vertices of a hypergraph considered at the moment. By identifying each vertex $v_i \in \mathcal{X}(\mathcal{H})$ with a variable $x_i \in R$, we may think of an edge E_i of a hypergraph as a monomial $x^{E_i} = \prod_{j \in E_i} x_j$ in R . Employing this idea, we may associate to every simple hypergraph $\mathcal{H}$, a squarefree monomial ideal in R . The **edge ideal** $I(\mathcal{H})$ of a hypergraph $\mathcal{H}$ is the ideal

$$I(\mathcal{H}) = (x^{E_i} ; E_i \in \mathcal{E}(\mathcal{H})) \subseteq R,$$

*Department of Mathematics, Stockholm University, 106 91 Stockholm, erice@math.su.se

generated “by the edges” of $\mathcal{H}$. This yields the **hypergraph algebra** $R/I(\mathcal{H})$.

In this way we obtain a 1-1 correspondence

$$\{\text{simple hypergraphs on } [n]\} \longleftrightarrow \{\text{squarefree monomial ideals } I \subseteq R = k[x_1, \dots, x_n]\}.$$

Recall that an **(abstract) simplicial complex** on vertex set $[n]$ is a collection, Δ , of subsets of $[n]$ with the property that $G \subseteq F$, $F \in \Delta \Rightarrow G \in \Delta$. The elements of Δ are called the **faces** of the complex and the maximal (under inclusion) faces are called **facets**. The **dimension**, $\dim F$, of a face F in Δ , is defined to be $|F| - 1$, and the dimension of Δ is defined as $\dim \Delta = \max\{\dim F; F \in \Delta\}$. Note that the empty set $\emptyset$ is the unique -1 dimensional face of every complex that is not the void complex $\{\}$ which has no faces. The dimension of the void complex may be defined as $-\infty$. The r -skeleton of a simplicial complex Δ , is the collection of faces of Δ of dimension at most r . Let $V \subseteq [n]$. We denote by Δ_V the simplicial complex

$$\Delta_V = \{F \subseteq [n]; F \in \Delta, F \subseteq V\}.$$

For convenience, we consider 0 to be a natural number, i.e., $\mathbb{N} = \{0, 1, 2, 3, \dots\}$. A vector $\mathbf{j} = (j_1, \dots, j_n) \in \{0, 1\}^n$ is called a squarefree vector in $\mathbb{N}^n$. We may identify $\mathbf{j}$ with the set $V \subseteq [n]$, where $i \in V$ precisely when $j_i = 1$. Since this correspondence between the V and the $\mathbf{j}$ is bijective, we may also denote Δ_V by $\Delta_{\mathbf{j}}$.

Given a simplicial complex Δ , we denote by $\mathcal{C}(\Delta)$ its reduced chain complex, and by $\tilde{H}_n(\Delta; k) = Z_n(\Delta)/B_n(\Delta)$ its n 'th reduced homology group with coefficients in the field k . In general we could use an arbitrary abelian group instead of k , but we will only consider the case when the coefficients lie in a field. For convenience, we define the homology of the void complex to be zero.

Let Δ be an arbitrary simplicial complex on $[n]$. The **Alexander dual simplicial complex** Δ^* to Δ , is defined by

$$\Delta^* = \{F \subseteq [n]; [n] \setminus F \notin \Delta\}.$$

Note that $(\Delta^*)^* = \Delta$.

The edge ideal was first introduced by R. H. Villarreal [16], in the case when $\mathcal{H} = \mathcal{G}$ is a simple graph. After that, hypergraph algebras has been widely studied. See for instance [4, 5, 9, 10, 11, 12, 13, 15, 17]. In [9], the authors use certain connectedness properties to determine a class of hypergraphs such that the hypergraph algebras have linear resolutions. Furthermore, nice recursive formulas for computing the Betti numbers are given.

Perhaps the most common way to study the connections between the combinatorial information contained in a hypergraph, and the algebraic information contained in the corresponding hypergraph algebra, is the one given by the *Stanley-Reisner correspondence*, which is a 1-1 correspondence:

$$\{\text{simplicial complexes on } [n]\} \longleftrightarrow \{\text{squarefree monomial ideals } I \subseteq R = k[x_1, \dots, x_n]\}$$

$$\Delta \rightsquigarrow I_\Delta.$$

Here, a monomial x^F is an element in I_Δ precisely when F is a non face in Δ . Note that using the above two 1-1 correspondences, we also get a 1-1 correspondence between the class of simple hypergraphs on $[n]$, and the class of simplicial complexes on $[n]$.

Let $\mathcal{H} = ([n], \mathcal{E}(\mathcal{H}))$ be a simple hypergraph and consider its edge ideal $I(\mathcal{H}) \subseteq R$. Note that $R/I(\mathcal{H})$ is precisely the Stanley-Reisner ring of the simplicial complex

$$\Delta(\mathcal{H}) = \{F \subseteq [n]; E \not\subseteq F, \forall E \in \mathcal{E}(\mathcal{H})\}.$$

This is called the **independence complex** of $\mathcal{H}$. The edges in $\mathcal{H}$ are precisely the minimal non faces of $\Delta(\mathcal{H})$.

Thus, we may think of the edges of a simple hypergraph as the minimal non faces of a simplicial complex or, equally well, the relations in the k -algebra $R/I(\mathcal{H})$. The connections between a (hyper)graph and its independence complex are explored in, for example [4, 7, 13].

Another way to use hypergraphs to investigate the properties of simplicial complexes was introduced in [5] by S. Faridi. Given a simplicial complex Δ , denote by $\{F_1, \dots, F_t\}$ the set of facets of Δ . Faridi then defines another squarefree monomial ideal, the **facet ideal** of Δ ,

$$\mathcal{F}(\Delta) = (x^F; F \text{ is a facet of } \Delta).$$

In several papers, for example [5, 6, 17], properties of simplicial complexes are studied via the combinatorial properties of their facet ideals. Note that the set of facets of Δ is a simple hypergraph.

In section 4, we introduce the classes of chordal and triangulated hypergraphs. The definition of triangulated hypergraph is almost identical to Definition 5.5 in [9], however, ours is more general. These classes of hypergraphs illustrates that d -uniform hypergraphs behaves much like ordinary simple graphs. However, there are familiar properties of graphs that do not translate immediately to d -uniform hypergraphs. See for instance Remark 4.1 and Example 1.

It is well known, see [8], that chordal graphs are characterized by the fact that they have perfect elimination orders. We show that this remain true for hypergraphs.

In Theorem 4.1 we show that the properties of being triangulated, chordal, and having a perfect elimination order, are equivalent also for hypergraphs.

In section 5 we introduce the class of generalized chordal hypergraphs, which includes the chordal hypergraphs, and show that the corresponding hypergraph algebras, $R/I(\mathcal{H})$, have linear resolutions. Our method of proof is a natural generalization of one used by R. Fröberg in [7]. There, Fröberg characterizes, in terms of the complementary graphs $\mathcal{G}^c$, precisely for what graphs $\mathcal{G}$ the graph algebras $R/I(\mathcal{G})$ have linear resolutions. Fröberg shows:

Theorem 1.1. *Let G be a simple graph on n vertices. Then $k[x_1, \dots, x_n]/I(\mathcal{G})$ has linear resolution precisely when $\mathcal{G}^c$ is chordal (rigid circuit, triangulated,...).*

By Theorem 5.1, we obtain a partial generalization of Fröberg's theorem.

Let Δ be an arbitrary simplicial complex, such that the Stanley-Reisner ring R/I_Δ has linear resolution. Then we know that the generators of I_Δ all have the same degree, d say. Thus, we may think of R/I_Δ as a hypergraph algebra $R/I(\mathcal{H})$ for some d -uniform hypergraph $\mathcal{H}$. However, we will look at things in another way. The **complementary hypergraph** $\mathcal{H}^c$, of a d -uniform hypergraph $\mathcal{H}$, is defined as the hypergraph on the same set of vertices as $\mathcal{H}$, and edge set

$$\mathcal{E}(\mathcal{H}^c) = \{F \subseteq \mathcal{X}(\mathcal{H}) ; |F| = d, F \notin \mathcal{E}(\mathcal{H})\}.$$

The edges of $\mathcal{H}^c$ may, in a natural way, be thought of as the $(d-1)$ -dimensional faces in the independence complex $\Delta(\mathcal{H})$, of $\mathcal{H}$. This is how Fröberg looks at things when he proves his theorem. We show that the complex $\Delta(\mathcal{H})$ is completely determined by the edges in $\mathcal{H}^c$, which gives us the notion of d -flag complexes.

2 Resolutions and Betti numbers

To every finitely generated graded module M over the polynomial ring $R = k[x_1, \dots, x_n]$, we may associate a **minimal (N-)graded free resolution**

$$0 \rightarrow \bigoplus_j R(-j)^{\beta_{l,j}(M)} \rightarrow \bigoplus_j R(-j)^{\beta_{l-1,j}(M)} \rightarrow \dots \rightarrow \bigoplus_j R(-j)^{\beta_{0,j}(M)} \rightarrow M \rightarrow 0$$

where $l \leq n$ and $R(-j)$ is the R -module obtained by shifting the degrees of R by j . Thus, $R(-j)$ is the graded R -module in which the grade i component $(R(-j))_i$ is R_{i-j} .

The natural number $\beta_{i,j}(M)$ is called the ij 'th **N-graded Betti number** of M . If M is multigraded we may equally well consider the $\mathbb{N}^n$ -graded minimal free resolution and Betti numbers of M . The difference lies just in the fact that we now use multigraded shifts $R(-\mathbf{j})$ instead of N-graded ones. The **total** i 'th Betti number is $\beta_i(M) = \sum_j \beta_{i,j}$. For further details on resolutions, graded rings and Betti numbers, we refer the reader to [1], sections 1.3 and 1.5.

The Betti numbers of M occur as the dimensions of certain vector spaces over $k = R/m$, where m is the unique maximal graded ideal in R . Accordingly, the Betti numbers in general depend on the characteristic of k .

A minimal free resolution of M is said to be **linear** if for $i > 0$, $\beta_{i,j}(M) = 0$ whenever $j \neq i + d - 1$ for some fixed natural number $d \geq 1$.

In connection to this we mention the *Eagon-Reiner theorem*.

Theorem 2.1. *Let Δ be a simplicial complex and Δ^* its Alexander dual complex. Then R/I_Δ is Cohen-Macaulay if and only if R/I_{Δ^*} has linear minimal free resolution.*

Proof. See [3], Theorem 3. □

3 Hochster's formula and the Mayer-Vietoris sequence

In topology one defines Betti numbers in a somewhat different manner. *Hochster's formula* provides a link between these and the Betti numbers defined above.

Theorem 3.1. (*Hochster's formula*). Let R/I_Δ be the Stanley-Reisner ring of a simplicial complex Δ . The non-zero Betti numbers of R/I_Δ are only in squarefree degrees $\mathbf{j}$ and may be expressed as

$$\beta_{i,\mathbf{j}}(R/I_\Delta) = \dim_k \tilde{H}_{|\mathbf{j}|-i-1}(\Delta_{\mathbf{j}}; k).$$

Hence the total i 'th Betti number may be expressed as

$$\beta_i(R/I_\Delta) = \sum_{V \subseteq [n]} \dim \tilde{H}_{|V|-i-1}(\Delta_V; k).$$

Proof. See [1], Theorem 5.5.1. □

If one has $\mathbb{N}^n$ -graded Betti numbers, it is easy to obtain the $\mathbb{N}$ -graded ones via

$$\beta_{i,j}(R/I_\Delta) = \sum_{\substack{\mathbf{j}' \in \mathbb{N}^n \\ |\mathbf{j}'|=j}} \beta_{i,\mathbf{j}'}(R/I_\Delta).$$

Thus,

$$\beta_{i,j}(R/I_\Delta) = \sum_{\substack{V \subseteq [n] \\ |V|=j}} \dim \tilde{H}_{|V|-i-1}(\Delta_V; k).$$

Recall that if we have an exact sequence of complexes,¹

$$\mathbf{0} \rightarrow \mathbf{L} \rightarrow \mathbf{M} \rightarrow \mathbf{N} \rightarrow \mathbf{0}$$

there is a long exact (reduced) homology sequence associated to it

$$\cdots \rightarrow H_r(N) \rightarrow H_{r-1}(L) \rightarrow H_{r-1}(M) \rightarrow H_{r-1}(N) \rightarrow \cdots.$$

When we prove Theorem 5.1, we will use this homology sequence in the special case where it is associated to a simplicial complex as follows.

Suppose we have a simplicial complex N and two subcomplexes L and M , such that $N = L \cup M$. This gives us an exact sequence of (reduced) chain complexes

$$0 \rightarrow \mathcal{C}.(L \cap M) \rightarrow \mathcal{C}.(L) \oplus \mathcal{C}.(M) \rightarrow \mathcal{C}.(N) \rightarrow 0.$$

The non trivial maps here are defined by $x \mapsto (x, -x)$ and $(x, y) \mapsto x + y$.

The long exact (reduced) homology sequence associated to this particular sequence, is called the Mayer-Vietoris sequence. More about the Mayer-Vietoris sequence can be found in [14], section 4.4.

¹That is, complexes of modules over some ring R .

4 The classes of chordal and triangulated hypergraphs

In this section, all hypergraphs are assumed to be simple and d -uniform.

Definition 4.1. Two distinct vertices x, y of a hypergraph $\mathcal{H}$ are **neighbours** if there is an edge $E \in \mathcal{E}(\mathcal{H})$, such that $x, y \in E$. For any vertex $x \in \mathcal{X}(\mathcal{H})$, the **neighbourhood of x** , denoted $N(x)$, is the set

$$N(x) = \{y \in \mathcal{X}(\mathcal{H}) ; y \text{ is a neighbour of } x\}.$$

If $N(x) = \emptyset$, x is called isolated. Furthermore, we let $N[x] = N(x) \cup \{x\}$ denote the **closed neighbourhood of x** .

Remark 4.1. Let $\mathcal{H}$ be a hypergraph and $V \subseteq \mathcal{X}(\mathcal{H})$. Denote by $N_V[x]$ the closed neighbourhood of x in the induced hypergraph $\mathcal{H}_V$. For ordinary graphs it is clear that $N_V[x] = N[x] \cap V$. This is not always the case for hypergraphs, as is shown in the example below. Note that the notation $N_V[x]$ will only occur in this remark and the example below. The fact that we do not make any greater use of it, is intimately connected to, and in a sense illustrates, the properties of the hypergraphs that we are to consider.

Example 1. Consider the hypergraph $\mathcal{H}$ on vertex set $\mathcal{X}(\mathcal{H}) = \{a, b, c, d, e\}$ and edge set $\mathcal{E}(\mathcal{H}) = \{\{a, b, c\}, \{a, d, e\}, \{b, c, d\}\}$. Let $V = \{a, b, c, d\}$. Then $N_V[a] = \{a, b, c\}$ but $N[a] \cap V = \{a, b, c, d\}$.

Recall the definition of the d -complete hypergraph:

Definition 4.2. The d -complete hypergraph, K_n^d , on a set of n vertices, is defined by

$$\mathcal{E}(K_n^d) = \binom{[n]}{d}$$

where $\binom{F}{d}$ denotes the set of all subsets of F , of cardinality d . If $n < d$, we interpret K_n^d as n isolated points.

If $\mathcal{H}$ is a hypergraph, we associate a simplicial complex $\Delta_{\mathcal{H}}$ to it in the following way:

Definition 4.3. Given a hypergraph $\mathcal{H} = (\mathcal{X}(\mathcal{H}), \mathcal{E}(\mathcal{H}))$, the **complex of $\mathcal{H}$** , $\Delta_{\mathcal{H}}$, is the simplicial complex

$$\Delta_{\mathcal{H}} = \{F \subseteq \mathcal{X}(\mathcal{H}) ; \binom{F}{d} \subseteq \mathcal{E}(\mathcal{H})\}$$

Note that this implies that if $F \subseteq \mathcal{X}(\mathcal{H})$, $|F| < d$, then $F \in \Delta_{\mathcal{H}}$.

Remark 4.2. $\Delta_{\mathcal{H}}$ is completely determined by $\mathcal{H}$. Indeed, all faces of dimension at least $d-1$ clearly is determined by $\mathcal{H}$, since each one correspond uniquely to a d -complete subhypergraph of $\mathcal{H}$.

Remark 4.3. Recall that a flag complex is a simplicial complex in which every minimal non face consists of precisely 2 elements. As one easily sees, such complex is determined by its 1-skeleton. According to the previous remark, **d -flag complexes**, i.e. complexes $\Delta_{\mathcal{H}}$ whose minimal non faces all have cardinality d , in a natural way generalizes flag complexes.

Proposition 4.1. $\Delta_{\mathcal{H}} = \Delta(\mathcal{H}^c)$, where $\Delta(\mathcal{H}^c)$ is the independence complex of $\mathcal{H}^c$.

Proof. The two complexes has the same set of vertices. $F \in \Delta(\mathcal{H}^c)$ precisely when $\binom{F}{d} \subseteq \mathcal{E}(\mathcal{H})$. Furthermore, $F \in \Delta(\mathcal{H}^c)$ for every $F \subseteq \mathcal{X}(\mathcal{H})$ with $|F| < d$. $\square$

Definition 4.4. Let Δ be a simplicial complex on a finite set, $\mathcal{X}$, of vertices. For any given $d \in \mathbb{N}$, the d -uniform hypergraph, $\mathcal{H}_d(\Delta)$, of Δ , is the hypergraph with vertex set $\mathcal{X}$, and with edge set

$$\mathcal{E}_d(\Delta) = \{F \in \Delta; |F| = d\}.$$

Proposition 4.2. Let $\mathcal{H}$ be a hypergraph and Δ an arbitrary d -flag complex on $\mathcal{X}(\mathcal{H})$. Then,

- $\mathcal{H}_d(\Delta_{\mathcal{H}}) = \mathcal{H}$
- $\Delta_{\mathcal{H}_d(\Delta)} = \Delta$

Proof. This follows directly from Definition 4.2 and Definition 4.3. $\square$

Definition 4.5. A hypergraph $\mathcal{H}$ is called **triangulated** if for every non empty subset $V \subseteq \mathcal{X}(\mathcal{H})$, either there exists a vertex $x \in V$ such that the induced hypergraph $\mathcal{H}_{N[x] \cap V}$ is isomorphic to a d -complete hypergraph K_n^d , $n \geq d$, or else the edge set of $\mathcal{H}_V$ is empty.

Definition 4.6. A hypergraph $\mathcal{H}$ is called **triangulated*** if for every non empty subset $V \subseteq \mathcal{X}(\mathcal{H})$, either there exists a vertex $x \in V$ such that $N[x] \cap V$ is a facet of $(\Delta_{\mathcal{H}})_V$ of dimension greater than or equal to $d - 1$, or else the edge set of $\mathcal{H}_V$ is empty.

Definition 4.7. A **chordal hypergraph** is a d -uniform hypergraph, obtained inductively as follows:

- K_n^d is a chordal hypergraph, $n, d \in \mathbb{N}$.
- If $\mathcal{G}$ is chordal, then so is $\mathcal{H} = \mathcal{G} \cup_{K_j^d} K_i^d$, for $0 \leq j < i$. (This we think of as glueing K_i^d to $\mathcal{G}$ by identifying some edges, or parts of some edges, of K_i^d with the corresponding part, K_j^d , of $\mathcal{G}$.)

Remark 4.4. For $d = 2$ this specializes precisely to the class of generalized trees, i.e. generalized n -trees for some n , as defined in [7].

Remark 4.5. In the special case of simple graphs, Definition 4.5 specializes precisely to the ordinary chordal (rigid circuit) graphs. Recall that a simple graph is called chordal if every induced cycle of length > 3 , has a chord. By considering minimal cycles, it is clear that a graph that is triangulated according to Definition 4.5, is chordal. Assume a graph $\mathcal{G}$ is chordal. It follows from Theorems 1 and 2 in [2], that the chordal graphs are precisely the generalized trees (see Remark 4.4). In a generalized tree we may easily find a vertex x , with the property that $\mathcal{G}_{N[x]}$ is complete, as follows: We know that $\mathcal{G} = \mathcal{G}' \cup_{K_j} K_i$, $0 \leq j < i$. Then, we just pick a vertex $x \in \mathcal{X}(K_i) \setminus \mathcal{X}(\mathcal{G}')$, since such x clearly has the property that $\mathcal{G}_{N[x]}$ is complete. Since every induced subgraph of a chordal graph is chordal, the same thing holds for every $\mathcal{G}_V$, $V \subseteq \mathcal{X}(\mathcal{G})$.

Another characterization of chordal graphs may be found in [8]. There it is shown that a simple graph is chordal precisely when it has a perfect elimination order. Recall that a perfect elimination order of a graph $\mathcal{G} = (\mathcal{X}, \mathcal{E})$ is an ordering of its vertices, $x_1 < x_2 < \dots < x_n$, such that for each i , $\mathcal{G}_{N[x_i] \cap \{x_i, x_{i+1}, \dots, x_n\}}$ is a complete graph. The concept of perfect elimination order is well suited for generalizations. We make the following

Definition 4.8. A hypergraph $\mathcal{H}$ is said to have a **perfect elimination order** if its vertices can be ordered $x_1 < x_2 < \dots < x_n$, such that for each i , either $\mathcal{H}_{N[x_i] \cap \{x_i, x_{i+1}, \dots, x_n\}}$ is isomorphic to a d -complete hypergraph K_n^d , $n \geq d$, or else x_i is isolated in $\mathcal{H}_{\{x_i, x_{i+1}, \dots, x_n\}}$

Note that this specializes precisely to the definition of perfect elimination order for simple graphs if we put $d = 2$.

Lemma 4.1. Let $\mathcal{H}$ be a hypergraph and $x \in V \subseteq \mathcal{X}(\mathcal{H})$ a vertex such that $\mathcal{H}_{N[x]} \cong K_m^d$, $m \geq d$. Then $\mathcal{H}_{N[x] \cap V}$ either is isomorphic to a d -complete hypergraph $K_{m'}^d$, $m' \geq d$, or else x is isolated in V .

Proof. Either $|N[x] \cap V| \geq d$ or else $|N[x] \cap V| < d$. □

Remark 4.6. The above lemma in some sense explains what goes on in the proofs hereafter. It also casts some light on the last comment made in Remark 4.1.

Lemma 4.2. If a hypergraph $\mathcal{H}$ with $\mathcal{E}(\mathcal{H}) \neq \emptyset$ has a perfect elimination order, then it has a perfect elimination order $x_1 < x_2 < \dots < x_n$ in which x_1 is not isolated.

Proof. Let $x_1 < x_2 < \dots < x_n$ be a perfect elimination order of $\mathcal{H}$, and put

$$t = \min\{i; x_i \text{ is not isolated}\}.$$

We claim that $x_t < \dots < x_n < x_1 < \dots < x_{t-1}$ also is a perfect elimination order of $\mathcal{H}$. Since $x_1, \dots, x_{t-1}$ are isolated, we need only verify that $\mathcal{H}_{N[x_i] \cap \{x_i, x_{i+1}, \dots, x_n, x_1, \dots, x_{t-1}\}} \cong K_{m_i}^d$ for some $m_i \geq d$, $i = t, \dots, n$. However, this is clear since $\mathcal{H}_{N[x_i] \cap \{x_i, x_{i+1}, \dots, x_n, x_1, \dots, x_{t-1}\}} = \mathcal{H}_{N[x_i] \cap \{x_i, x_{i+1}, \dots, x_n\}}$. □

Lemma 4.3. If a hypergraph $\mathcal{H}$ is triangulated (triangulated*, chordal), or, has a perfect elimination order, then so does $\mathcal{H}_V$ for every $V \subseteq \mathcal{X}(\mathcal{H})$.

Proof. Let $V \subseteq \mathcal{X}(\mathcal{H})$. If $\mathcal{E}(\mathcal{H}_V) = \emptyset$, $\mathcal{H}_V$ clearly is triangulated and triangulated*. It is also chordal since we can add one vertex at a time until we have the desired discrete hypergraph, and any ordering of V yields a perfect elimination order. Thus we may assume that $\mathcal{E}(\mathcal{H}_V) \neq \emptyset$.

The lemma is clear for the classes of triangulated and triangulated* hypergraphs, since if $W \subseteq V$, we have that $(\mathcal{H}_V)_W = \mathcal{H}_W$. Now, let $\mathcal{H} = \mathcal{G} \bigcup_{K_j^d} K_i^d$, $0 \leq j < i$, be chordal. If $V \subseteq \mathcal{X}(\mathcal{G})$, or if $V \subseteq \mathcal{X}(K_i^d)$, we are done by induction. If this is not the case, it is easy to realize that $\mathcal{H}_V = \mathcal{G}_V \bigcup_{(K_j^d)_V} (K_i^d)_V$. Since $\mathcal{G}_V$ is chordal by induction, the result follows. Finally, assume $\mathcal{H}$ has a perfect elimination order $x_1 < x_2 < \dots < x_n$. Then V inherits an

ordering $x_{i_1} < x_{i_2} < \dots < x_{i_{|V|}}$. The fact that this is a perfect elimination order of $\mathcal{H}_V$ follows from Lemma 4.1. $\square$

Theorem 4.1. *Let $\mathcal{H} = (\mathcal{X}(\mathcal{H}), \mathcal{E}(\mathcal{H}))$ be a d -uniform hypergraph. Then the following are equivalent.*

- (i) $\mathcal{H}$ is triangulated.
- (ii) $\mathcal{H}$ is triangulated*.
- (iii) $\mathcal{H}$ is chordal.
- (iv) $\mathcal{H}$ has a perfect elimination order.

Proof. Due to Lemma 4.3, we need only consider the full set $\mathcal{X}(\mathcal{H})$ of vertices in our arguments, and we may assume that $\mathcal{E}(\mathcal{H}) \neq \emptyset$.

(i) $\Rightarrow$ (ii). Since we assume $\mathcal{E}(\mathcal{H}) \neq \emptyset$ and consider only the case where $V = \mathcal{X}(\mathcal{H})$, there is a vertex x such that $\mathcal{H}_{N[x]} \cong K_n^d$, $n \geq d$. Then, $N[x]$ clearly is a face in $\Delta_{\mathcal{H}}$ of dimension at least $d - 1$. Furthermore it has to be a facet, since if there were a $y \in \mathcal{X}(\mathcal{H})$, $y \neq x$, such that $N[x] \cup \{y\} \in \Delta_{\mathcal{H}}$, then there would exist an edge E with $x, y \in E$. Hence, $y \in N[x]$.

(ii) $\Rightarrow$ (i). By assumption, there is a vertex x such that $N[x]$ is a facet in $\Delta_{\mathcal{H}}$ of dimension greater than or equal to $d - 1$, whence it is clear (from the definition of $\Delta_{\mathcal{H}}$) that $\mathcal{H}_{N[x]} \cong K_n^d$ for some $n \geq d$.

(i) $\Rightarrow$ (iii). By assumption there is a vertex $x \in \mathcal{X}(\mathcal{H})$ such that $\mathcal{H}_{N[x]} \cong K_n^d$, for some $n \geq d$. Let $\mathcal{G}$ be the induced hypergraph on $\mathcal{X}(\mathcal{H}) \setminus \{x\}$. Then $\mathcal{E}(\mathcal{G})$ consists of all edges of $\mathcal{H}$, except those that contain x . This yields $\mathcal{H} = \mathcal{G} \cup_{\mathcal{K}} K_n^d$, where $\mathcal{K} = K_{|N(x)|}^d$ on vertex set $N(x)$, and by induction we are done.

(iii) $\Rightarrow$ (i). Assume $\mathcal{H} = \mathcal{G} \cup_{K_j^d} K_i^d$, $0 \leq j < i$, is chordal, where $\mathcal{G}$ is chordal by construction. If $i \geq d$, any vertex $x \in \mathcal{X}(K_i^d) \setminus \mathcal{X}(\mathcal{G})$ will do, since $\mathcal{H}_{N[x]} \cong K_i^d$ for such x . If $i < d$, we find, by induction, a vertex $x \in \mathcal{X}(\mathcal{G})$ with the property that $\mathcal{H}_{N[x]} = \mathcal{G}_{N[x]} \cong K_n^d$ for some $n \geq d$, since otherwise the edge set of $\mathcal{H}$ would be empty, contrary to our assumptions.

(i) $\Rightarrow$ (iv). By assumption we find a vertex $x = x_1$ such that $\mathcal{H}_{N[x_1]} \cong K_n^d$, $n \geq d$. Since the induced hypergraph on $\mathcal{X}(\mathcal{H}) \setminus \{x_1\}$ is triangulated, by induction it has a perfect elimination order $x_2 < \dots < x_n$. If we put $x_1 < x_2$ we are done.

(iv) $\Rightarrow$ (i). By Lemma 4.2 there is a perfect elimination order $x_1 < \dots < x_n$, such that $\mathcal{H}_{N[x_1] \cap V} \cong K_m^d$ for some $m \geq d$. $\square$

4.1 Some examples

In [4], we considered hypergraph generalizations of the well known complete and complete multipartite graphs. We use these to create some examples of chordal hypergraphs.

Recall from [4] the definition of the d -complete bipartite hypergraph $K_{n,m}^d$: This is the hypergraph on a vertex set that is a disjoint union, $[n] \sqcup [m]$, of two finite sets. The edge set consists of all sets $V \subseteq [n] \sqcup [m]$, $|V| = d$, such that $V \cap [n] \neq \emptyset \neq V \cap [m]$.

Example 2. Here we consider the complement $\mathcal{H} = (K_{n,m}^d)^c$ of $K_{n,m}^d$. We claim that $\mathcal{H}$ is chordal. It is easy to see, considering the relations in the Stanley-Reisner ring, that $\Delta_{\mathcal{H}}$ looks like

$$(\Delta_n \sqcup \Delta_m) \cup \Gamma_{d-2}([n] \cup [m])$$

where Δ_r is the full simplex on $[r]$, and $\Gamma_{d-2}([n] \cup [m])$ is the $(d-2)$ -skeleton of the full simplex on $[n] \sqcup [m]$.

Clearly, the d -uniform hypergraph of this complex, in other words $\mathcal{H}$, is the disjoint union two d -complete hypergraphs,

$$\mathcal{H} = K_n^d \cup_{K_0^d} K_m^d,$$

so $\mathcal{H}$ is chordal.

Example 3. Now consider the complex $\Delta_{K_{n,m}^d}$, of $K_{n,m}^d$. If $n, m < d$, we have an isomorphism $K_{n,m}^d \cong K_{n+m}^d$, so in this case $K_{n,m}^d$ is chordal. If n or m is greater than or equal to d , $K_{n,m}^d$ is not chordal. This is because no matter which vertex x we choose, the induced hypergraph on $N[x]$ cannot be d -complete, since it would then contain an edge lying entirely in either $[n]$ or $[m]$, which is impossible.

The general case of the d -complete multipartite hypergraph, $K_{n_1, \dots, n_t}^d$, is similar. $K_{n_1, \dots, n_t}^d$ is chordal only when $n_i < d$ for every $i = 1, \dots, t$. The arguments are the same as in the bipartite case.


Another kind of complete hypergraph, is the $d(a, b)$ -complete hypergraph $\mathcal{H} = K_{n,m}^{d(a,b)}$, where $d = a + b$, $a, b \geq 1$. Here $\mathcal{X}(\mathcal{H}) = [n] \sqcup [m]$, and $\mathcal{E}(\mathcal{H}) = \binom{[n]}{a} \times \binom{[m]}{b}$.

Example 4. Consider the complex of $K_{n,m}^{d(a,b)}$. Pick any vertex x and consider $N[x]$. If the induced hypergraph $(K_{n,m}^{d(a,b)})_{N[x]}$ is to be complete, both n and m must be smaller than d , and at least one of the two equations $n = a$, $m = b$ must hold. Otherwise we obtain a contradiction since $K_{n,m}^{d(a,b)}$ would then contain an edge of the wrong shape. If n and m satisfy these conditions, the hypergraph is chordal.

5 Generalized chordal hypergraphs

It is easy to find an example of a d -uniform hypergraph $\mathcal{H}$ that is not chordal, but such that the Stanley-Reisner ring of $\Delta_{\mathcal{H}}$ has linear resolution.

Example 5: Let $\mathcal{H}$ be the 3-uniform hypergraph with $\mathcal{X}(\mathcal{H}) = \{a, b, c, d\}$, and $\mathcal{E}(\mathcal{H}) = \{\{a, b, c\}, \{a, c, d\}, \{a, b, d\}\}$. The following simple picture lets us visualize $\mathcal{H}$.



$R/I_{\Delta_{\mathcal{H}}}$ has linear resolution, but $\mathcal{H}$ is not chordal.

If Δ is a simplicial complex on $[n]$ and E is a finite set, we denote by $\Delta \cup E$ the simplicial complex on $[n] \cup E$ whose set of facets, $\mathcal{F}(\Delta \cup E)$, is $\mathcal{F}(\Delta) \cup \{E\}$. Similarly, if $\mathcal{H}$ is a (not necessarily d -uniform) hypergraph and E a finite set, we denote by $\mathcal{H} \cup E$ the hypergraph on $\mathcal{X}(\mathcal{H}) \cup E$ whose edge set is $\mathcal{E}(\mathcal{H} \cup E) = \mathcal{E}(\mathcal{H}) \cup \{E\}$.

Definition 5.1. A **generalized chordal hypergraph** is a d -uniform hypergraph, obtained inductively as follows:

- K_n^d is a generalized chordal hypergraph, $n, d \in \mathbb{N}$.
- If $\mathcal{G}$ is generalized chordal, then so is $\mathcal{H} = \mathcal{G} \cup_{K_j^d} K_i^d$, for $0 \leq j < i$.
- If $\mathcal{G}$ is generalized chordal and $E \subseteq \mathcal{X}(\mathcal{G})$ a finite set, $|E| = d$, such that at least one element of $\binom{E}{d-1}$ is not a subset of any edge of $\mathcal{G}$, then $\mathcal{G} \cup E$ is generalized chordal.

Remark 5.1. It is clear that every chordal hypergraph is also a generalized chordal hypergraph. Furthermore, for $d = 2$ chordal graphs and generalized chordal graphs are the same.

Theorem 5.1. Let $\mathcal{H} = (\mathcal{X}(\mathcal{H}), \mathcal{E}(\mathcal{H}))$ be a generalized chordal hypergraph and k a field of arbitrary characteristic. Then the Stanley-Reisner ring of $\Delta_{\mathcal{H}}$ has linear resolution.

Proof. We consider the three instances of Definition 5.1 one at a time. If $\mathcal{H} \cong K_n^d$ we are done, since if $n \geq d$ we have a simplex so the situation is trivial, and if $n < d$ the claim is proved for example in [4], Theorem 3.1. So, we may assume $\mathcal{H} \not\cong K_n^d$. Let $\mathcal{H} = \mathcal{G} \cup_{K_j^d} K_i^d$, $0 \leq j < i$, where $\mathcal{G}$ is generalized chordal. Let C and B be the simplices determined by K_j^d and K_i^d , respectively, and consider the complex $\Delta'_{\mathcal{H}} = \Delta_{\mathcal{G}} \cup B$. Note that $B \cap \Delta_{\mathcal{G}} = C$, $B \neq C$. We first show that $\Delta'_{\mathcal{H}}$ has linear resolution. For every $V \subseteq \mathcal{X}(\mathcal{H})$, we have an exact sequence of chain complexes

$$0 \rightarrow \mathcal{C}.(C_V) \rightarrow \mathcal{C}..(\Delta_{\mathcal{G}})_V) \oplus \mathcal{C}..(B_V) \rightarrow \mathcal{C}..(\Delta'_{\mathcal{H}})_V) \rightarrow 0.$$

By induction, via Hochster's formula, we know that $(\Delta_{\mathcal{G}})_V$ can have non zero homology only in degree $d - 2$. But then, since both B_V and C_V are simplices and accordingly have no homology at all, by considering the Mayer-Vietoris sequence we conclude that the only possible non zero homologies of $(\Delta'_{\mathcal{H}})_V$ lies in degree $d - 2$.

Note that it is not in general true that $\Delta_{\mathcal{H}} = \Delta'_{\mathcal{H}}$. In fact, this holds only when $d = 2$.

However, the difference between the two complexes is easy to understand, and we may use the somewhat easier looking $\Delta'_{\mathcal{H}}$ to show that $\Delta_{\mathcal{H}}$ has linear resolution as well.

To this end, let $\Gamma_{d-2}(\mathcal{X}(\mathcal{H}))$ be the $(d-2)$ -skeleton of the full simplex on vertex set $\mathcal{X}(\mathcal{H})$. Then one sees that

$$\Delta_{\mathcal{H}} = \Delta'_{\mathcal{H}} \cup \Gamma_{d-2}(\mathcal{X}(\mathcal{H})).$$

The $(d-2)$ -faces that we add to $\Delta'_{\mathcal{H}}$ to obtain $\Delta_{\mathcal{H}}$, can certainly not cause any homology in degrees greater than $d-2$, that did not already exist in $\Delta'_{\mathcal{H}}$. Indeed, suppose $\sum_i a_i \sigma_i$ is a cycle in a degree $r > d-2$, where $a_i \in k$ and the σ_i 's are faces of $\Delta_{\mathcal{H}}$, of dimension r . Since every face σ_i actually lies in $\Delta'_{\mathcal{H}}$, it follows that $\sum_i a_i \sigma_i$ is a cycle also in $\Delta'_{\mathcal{H}}$. Thus, if $\Delta'_{\mathcal{H}}$ has linear resolution, so does $\Delta_{\mathcal{H}}$.

Finally, let $\mathcal{H} = \mathcal{G} \cup E$. Let $F_1, \dots, F_t$ be the elements of $\binom{E}{d-1}$ that are not subsets of any edge of $\mathcal{G}$. Note that $\Delta_{\mathcal{H}} = \Delta_{\mathcal{G}} \cup E$. Take $V \subseteq \mathcal{X}(\mathcal{H})$. If $E \not\subseteq V$, then $(\Delta_{\mathcal{H}})_V = (\Delta_{\mathcal{G}})_V$, so, by induction we conclude that the only possible non zero homologies of $(\Delta_{\mathcal{H}})_V$ lies in degree $d-2$. Hence we may assume that $E \subseteq V$. Then we have an exact sequence

$$0 \rightarrow \mathcal{C}.((\Delta_{\mathcal{G}} \cap E)_V) \rightarrow \mathcal{C}.((\Delta_{\mathcal{G}})_V) \oplus \mathcal{C}.(E_V) \rightarrow \mathcal{C}.((\Delta_{\mathcal{H}})_V) \rightarrow 0.$$

Note that E_V is a simplex so it has no homology, and, by induction, we know that $R/I_{\Delta_{\mathcal{G}}}$ has linear resolution. Using Hochster's formula, we may conclude that $\tilde{H}_{d-1}((\Delta_{\mathcal{G}})_V; k) = 0$. Hence, the Mayer-Vietoris sequence obtained from the above exact sequence looks as follows:

$$0 \rightarrow \tilde{H}_{d-1}((\Delta_{\mathcal{H}})_V) \rightarrow \tilde{H}_{d-2}((\Delta_{\mathcal{G}} \cap E)_V) \rightarrow \tilde{H}_{d-2}((\Delta_{\mathcal{G}})_V) \rightarrow \tilde{H}_{d-2}((\Delta_{\mathcal{H}})_V) \rightarrow 0.$$

Let $z = \sum_j a_j \sigma_j$ be an element in $Z_{d-1}((\Delta_{\mathcal{H}})_V)$, where $\sigma_1 = E$. Consider the expression for the derivative of this cycle

$$0 = d(z) = \dots + \sum_{i=1}^t \pm a_i F_i + \dots.$$

Since $\sum_{i=1}^t \pm a_i F_i$ only can come from $d(E)$, we conclude that $a_1 = 0$. Hence $z \in Z_{d-1}((\Delta_{\mathcal{G}})_V)$, and, using Hochster's formula, we may conclude that the Stanley-Reisner ring of $\Delta_{\mathcal{H}}$ has linear resolution. □

Corollary 5.1. *Let $\mathcal{H} = (\mathcal{X}(\mathcal{H}), \mathcal{E}(\mathcal{H}))$ be a generalized chordal hypergraph and k a field of arbitrary characteristic. Then the Stanley-Reisner ring $R/I_{\Delta_{\mathcal{H}}}^*$ of the Alexander dual complex $\Delta_{\mathcal{H}}^*$ is Cohen-Macaulay.*

Proof. This follows by the Eagon-Reiner theorem. □

Corollary 5.2. *Theorem 5.1 and Corollary 5.1 in particular applies to triangulated and triangulated* hypergraphs, and also to hypergraphs that have perfect elimination orders.*

Question 1: Is there a hypergraph $\mathcal{H}$ such that the Stanley-Reisner ring of $\Delta_{\mathcal{H}}$ has linear resolution over any field k , but that is not a generalized chordal hypergraph?

Question 2: If $\mathcal{H}$ is a generalized chordal hypergraph, are there more equivalent characterizations of $\mathcal{H}$ similar to those for a chordal hypergraph given in Theorem 4.1?

References

- [1] W. Bruns and J. Herzog. *Cohen-Macaulay rings*. Cambridge University Press, revised edition, 1998.
- [2] G. A. Dirac. On rigid circuit graphs. *Abh. Math. Sem. Univ. Hamburg*, 25:71–76, 1961.
- [3] J. A. Eagon and V. Reiner. Resolutions of Stanley-Reisner rings and Alexander Duality. *J. Pure Appl. Algebra*, 130:265–275, 1998.
- [4] E. Emtander. Betti numbers of hypergraphs. *arXiv:0711.3368*. To appear in *Communications in algebra*.
- [5] S. Faridi. The facet ideal of a simplicial complex. *Manuscripta Math.*, 242:92–108, 2002.
- [6] S. Faridi. Cohen-Macaulay Properties of Square-Free Monomial Ideals. *J. Combin. Theory*, Ser. A 109:299–329, 2005.
- [7] R. Fröberg. On Stanley-Reisner rings. *Topics in Algebra*, 26, 1990.
- [8] D. R. Fulkerson and O. A. Gross. Incidence matrices and interval graphs. *Pacific J. Math.*, 15:835–855, 1965.
- [9] H. T. Hà and A. Van Tuyl. Monomial ideals, edge ideals of hypergraphs, and their graded Betti numbers. *arXiv:math/0606539*, 2006.
- [10] H. T. Hà and A. Van Tuyl. Resolutions of square-free monomial ideals via facet ideals: a survey. *arXiv:math/0604301*, 2006.
- [11] H. T. Hà and A. Van Tuyl. Splittable ideals and resolutions of monomial ideals. *J. Algebra*, 309(1):405–425, 2007.
- [12] J. Herzog and M. Kühl. On the Betti numbers of finite pure and linear resolutions. *Comm. Algebra*, 12:1627–1646, 1984.
- [13] S. Jacques. *Betti Numbers of Graph Ideals*. PhD thesis, University of Sheffield, 2004. *arXiv:math/0410107*.
- [14] C. R. F. Maunder. *Algebraic Topology*. Dover publications Inc., New York, 1996.
- [15] S. Morey, E. Reyes, and R. H. Villarreal. Cohen-Macaulay, Shellable and unmixed clutters with a perfect matching of Knig type. *arXiv:0708.3111v3*, 2007.
- [16] R. H. Villarreal. Cohen-Macaulay graphs. *Manuscripta Math.*, 66:277–293, 1990.
- [17] X. Zheng. Resolutions of facet ideals. *Comm. Algebra*, 32(6):2301–2324, 2004.